

## **Effects of Hot Boning and Postmortem Aging Time on Camel Meat Tenderness**

**A. N. Al-Owaimer**

*Department of Animal Production, College of Agriculture,  
King Saud University, P. O. Box 2460, Riyadh 11451, Saudi Arabia.*

(Received 13/10/1421; accepted for publication 18/12/1421)

**Abstract.** An experiment was conducted to determine the effects of hot boning and postmortem aging time on camel meat tenderness. Six male camels (average carcass weight 133 kg) were used. Longissimus muscles were excised at 1 or 24 h postmortem and aged at 2 °C for 1, 3, 7, 10, 14 d postmortem. At the end of each period, shear force, myofibril fragmentation index (MFI), sarcomere length, drip and cooking losses were determined. Hot boning significantly ( $p < 0.05$ ) increased shear force at day 3 and 7 compared to intact muscle. Postmortem aging has reduced shear force during the first 7 days of postmortem. Most of the increases in myofibril fragmentation index (MFI) happened within the first 7 days of postmortem and MFI was reduced by hot boning compared to intact muscle. Sarcomere length decreased ( $p < 0.5$ ) by hot boning on day 3 and 7 compared to intact muscle. The results of this investigation indicate that hot boning had no effect on camel meat tenderness within the first 24-h postmortem, and camel meat tenderness can be improved by aging up to 7 days of postmortem.

### **Introduction**

Meat tenderness plays an important role for consumer satisfaction and meat price. Consumer satisfaction is the main goal for meat producers and retailers. One of the most important factors affecting consumer satisfaction is meat tenderness. There are different factors affecting meat tenderness, including animal age [1], sex [2], and muscle fiber size [3]. Also, postmortem aging time has been reported to improve meat tenderness in different animal species [4-6]. However, Koohmaraie [7] concluded that beef should be stored for 10-14 d, lamb for 10 d, and pork 5 d to get the maximum benefits of improving meat tenderness during postmortem storage.

The desert Arabian camel (*Camelus dromedarius*) constitutes 17 million heads in the world [8]. In certain areas, camel meat replaced beef and mutton [9]. Camel meat preference comes third [10] and fifth [11] meat of choice among Saudi Arabian citizen.

Several studies have been published concerning the physical characteristics, chemical composition, sensory properties and nutritive values of Najdi camel meat [12-15]. Informations relating to the effect of hot boning and postmortem aging time on camel meat tenderness are needed. Therefore, the objective of the present study is to investigate the effect of hot boning and aging time on the tenderness of camel longissimus muscle.

### **Materials and Methods**

Six Najdi male camels of similar background were slaughtered at Labone farm (Al-Kharj, Saudi Arabia), adressed and weighed. Within 1 h postmortem, the longissimus dorsi muscle (LM) of the right side was hot-boned from each carcass between the 7<sup>th</sup> and 11<sup>th</sup> rib region, whereas, the left side was intact and both were shipped to Meat Lab at King Saud University, Riyadh. All hot-boned muscles and intact sides were chilled for 24 h at 2 °C, thereafter, the LM of the intact side was separated from the bone. The muscles were cut into one inch steaks, vacuum-packaged using Turbovac vacuum machine (Model SB45H, hertogenbosch, the Netherlands), and stored at 2 °C for 3, 7, 10, and 14 days. With the exception of drip loss, all measurements were made at 1, 3, 7, 10 and 14 days postmortem for both hot boning and intact muscles.

#### **Drip loss**

At 3, 7, 10 and 14 days postmortem, meat samples were removed from the vacuum packages and purge were weighed and percentage of drip loss were calculated.

#### **Cooking loss**

Two steaks from each hot boned and intact muscles were weighed and cooked on Teval superbarbecue electric broiler (Teval, France) to an internal temperature of 70°C. Steaks were turned over after reaching 40°C. Internal temperature was monitored by Fisher alarm thermometer (Model No. 15-077-8B, Fisher Scientific, USA). After cooking, steaks were allowed to equilibrate to room temperature (23°C), then weighed and percentage of cooking loss was calculated.

#### **Shear force**

Steaks used for cooking loss experiment were used. Six 1.27 cm diameter cores from each steak were removed parallel to the long axis of the muscle fibers, then the cores were sheared perpendicular to the long axis of the fibers using the Warner-Bratzler shear force instrument, and the average value were recorded in kilogram/cm<sup>2</sup>.

#### **Myofibril fragmentation index**

At 1, 3, 7, 10, 14 d postmortem, myofibril fragmentation index (MFI) were determined on fresh muscle samples according to the procedure of Culler *et al.* [16].

### Sarcomere length

Three myofibrils were taken from each cut using forceps and surgical plate. The sarcomere lengths of the myofibrils were determined using the Laser head (Model 2.0MW) according to the method of Cross *et al.* [17].

### Statistical analysis

Data were analyzed by split plot design using GLM procedure of SAS [18]. Animal represented the whole plot while, hot boning and intact muscles defined the split plot. Treatment effect included five aging times. The means were considered significant at  $P$  level of  $<0.05$ .

## Result and Discussion

### Cooking and drip losses

The effect of hot boning and aging time on cooking and drip losses are presented in Table 1. Storage time and hot boning had little effect on cooking and drip losses. Parrish *et al.* [31] found that until 7 days postmortem storage time, there were no effect on cooking loss. Cooking characteristics were not affected by postmortem aging and hot-boned treatment [20, 22]. Conflicting study was reported by Klont *et al.* [29] who noted that deboning at 24 h after slaughter had led to significantly higher drip loss at 3, 7, and 14 d postmortem compared to 48 h deboning.

### Shear force

The Warner-Bratzler shear force results revealed that hot-boned samples had higher ( $P < 0.05$ ) shear force values on days 3 and 7 compared to the intact samples (Table 1). On day one, hot boning had no effect on shear force of camel longissimus muscle compared to intact muscle and this is beneficial to camel meat industry and Saudi consumers, who mostly consume camel meat within 24 h of postmortem. This result was in agreement with Wheeler *et al.* [19] who noted that hot boning had no effect ( $P > 0.05$ ) on shear force of beef biceps femoris muscle. Schmidt and Keman [20] found that no differences in any palatability traits of longissimus muscle biceps femoris or semimembranosus muscle that were hot-boned at 1 h postmortem compared with conventional process cuts.

The effect of postmortem aging time on shear force was presented in Table 1. Shear force value decreased gradually as the postmortem aging time increased in both hot boning and intact samples. There was no significant difference on tenderness after 7 and 10 days for the intact and hot boned samples, respectively. These results were in agreement with study of Koochmorie *et al.* [21] who reported that as the postmortem storage time increased, shear force values decreased for beef, lamb and pig and the majority of the improvement in tenderness occurred within 7 d of aging postmortem time. However, increasing aging time far over 7 days has no improvement in beef tenderness [22-24].

**Table 1. Effects of hot boning and postmortem aging time on camel meat traits**

Measurements	Postmortem (day)	Hot-boned	Intact
Cooking loss	1	28.00 <sup>a</sup> <sub>w</sub>	35.53 <sup>b</sup> <sub>w</sub>
	3	30.04 <sup>a</sup> <sub>w</sub>	29.71 <sup>a</sup> <sub>y</sub>
	7	37.90 <sup>a</sup> <sub>yx</sub>	37.70 <sup>a</sup> <sub>w</sub>
	10	35.03 <sup>a</sup> <sub>yz</sub>	36.60 <sup>a</sup> <sub>w</sub>
	14	31.05 <sup>a</sup> <sub>wxz</sub>	39.00 <sup>b</sup> <sub>w</sub>
	Drip loss	1	ND
3		3.50 <sup>a</sup> <sub>w</sub>	3.38 <sup>a</sup> <sub>w</sub>
7		6.50 <sup>a</sup> <sub>w</sub>	4.52 <sup>a</sup> <sub>w</sub>
10		6.26 <sup>a</sup> <sub>w</sub>	9.57 <sup>a</sup> <sub>y</sub>
14		5.12 <sup>a</sup> <sub>w</sub>	9.50 <sup>b</sup> <sub>y</sub>
Shear force (Kg/cm <sup>2</sup> )		1	4.02 <sup>a</sup> <sub>w</sub>
	3	4.36 <sup>a</sup> <sub>w</sub>	3.47 <sup>b</sup> <sub>wx</sub>
	7	3.75 <sup>a</sup> <sub>wx</sub>	2.98 <sup>b</sup> <sub>yx</sub>
	10	3.03 <sup>a</sup> <sub>yx</sub>	3.01 <sup>a</sup> <sub>wx</sub>
	14	2.81 <sup>a</sup> <sub>y</sub>	2.90 <sup>a</sup> <sub>yx</sub>
	MFI	1	25.21 <sup>a</sup> <sub>w</sub>
3		36.52 <sup>a</sup> <sub>y</sub>	51.30 <sup>b</sup> <sub>y</sub>
7		62.93 <sup>a</sup> <sub>x</sub>	79.05 <sup>b</sup> <sub>x</sub>
10		86.30 <sup>a</sup> <sub>z</sub>	83.02 <sup>a</sup> <sub>wz</sub>
14		90.02 <sup>a</sup> <sub>z</sub>	91.01 <sup>a</sup> <sub>z</sub>
Sarcomere length (μm)		1	1.72 <sup>a</sup> <sub>w</sub>
	3	1.34 <sup>a</sup> <sub>y</sub>	1.63 <sup>b</sup> <sub>w</sub>
	7	1.51 <sup>a</sup> <sub>yz</sub>	1.70 <sup>a</sup> <sub>w</sub>
	10	1.57 <sup>a</sup> <sub>wxz</sub>	1.73 <sup>b</sup> <sub>w</sub>
	14	1.48 <sup>a</sup> <sub>xyz</sub>	1.71 <sup>b</sup> <sub>w</sub>

<sup>ab</sup> means in the same row followed by different superscripts are significantly different ( $p < 0.05$ ).

<sup>w-z</sup> means in the same column followed by different superscripts are significantly different ( $p < 0.05$ ).

ND = Not determined.

N = 6.

The little change in shear force after 7 and 10 day of postmortem may be due to the connective tissue component rather than myofibril components [25]. The response of connective tissue to aging is generally theorized to be less than that of myofibril component [26].

### Myofibril fragmentation index MFI

The effect of hot boning and postmortem aging time on MFI of camel meat are presented in Table 1. MFI increased progressively during the first 7 days postmortem, however, 58% and 80% of the changes in MFI values occurred during the first 7 day postmortem for hot boning and intact muscles, respectively. Little changes have been

observed for MFI from day 7 to day 14. The changes in MFI have been reported to relate to the degradation of the myofibrillar protein during postmortem storage [27]. Taylor *et al.* [23] found that 46% of the increase in MFI had occurred by 3 d postmortem. Wheeler *et al.* [28] reported that MFI significantly increased after 3 d postmortem storage. Other study by Koohmaraie *et al.* [23] monitor postmortem changes in MFI begin immediately after slaughter and continuing until 7 and 14 d postmortem and found that 61 to 64% of the increase in MFI had occurred by 3 days of postmortem compared with 7 or 14 d. In this study, the parallel increases in MFI values and decrease shear value from day 1 to day 7 are the most dramatic and the change from day 7 to day 14 in shear value are in turn reduced.

### Sarcomere length

Sarcomere length was decreased significantly ( $p < 0.05$ ) by hot boning from day 1 to day 3 (table 1); the decrease was 22.1% on day 3 compared to day 1. This result was supported by Klont *et al.* [29] who reported that sarcomere length was significantly ( $P < 0.05$ ) shorter for loin deboning at 24 h than for loin from the same carcass deboning at 48 h. Herring *et al.* [30] noted that sarcomere of bovine muscle length was shorter when excised from its attachment. Despite the small decrease in sarcomere length on day 3 for the intact sample, there was no significant change on day 3, 7, 10, and 14 postmortem compared to day 1. This result in agreement with the study of Parrish *et al.* [31] who noted that the improvement of bovine muscle did occur in tenderness between 1 and 3 days postmortem, but very little change occurred in sarcomere length. Improved tenderness during postmortem aging, therefore, seems related to the fragmentation of myofibrils than to the changes in sarcomere length. Wheeler *et al.* [32] concluded that the increases in tenderness during aging were due to proteolysis degradation not to the increases of sarcomere length. Other study reported that the increase in tenderness does not seem to parallel increases in sarcomere length [31].

Based on our results, we concluded that camel meat could be hot-boned and consumed within 24 h and the optimum aging time for improving camel meat tenderness was 7 days.

**Acknowledgment.** The author would like to thank Dr. Suliman Zahran and research assistants of animal production department for their help in this study.

### References

- [1] Smith, G.C., Cross, H.R., Carpenter, Z.L., Murphey, C.E., Savell, J. W., Abraham, H.C. and Davis, G.W. "Relationship of USDA Maturity Groups to Palatability of Cooked Beef". *J. Food Sci.*, 47 (1982), 1100-1107.
- [2] Albaugh, A., Carroll, F.D., Ellis, K.W. and Albaugh, R. "Comparison of Carcass and Meat from Steaks, Short Scrotum Bulls Intact Bulls". *J. Anim. Sci.*, 41 (1975), 1627-1631.
- [3] Seideman, S.C., Koohmaraie, M. and Crouse, J.D. "The Influence of Muscle Fiber Size on Tenderness in A-maturity Heifers". *J. Food Qual.*, 11 (1988), 27-34.
- [4] Field, R.A., Riley, M.C. and Chang, Y.O. "Free Amino Acid Changes in Different Aged Bovine Muscles and Their Relationship to Shear Values". *J. Food Sci.*, 36 (1971), 611-612.

- [5] Jennings, T.G., Berry, B.W. and Joseph, A.L. "Influence of Fat Thickness, Marbling and Length of Aging on Beef Palatability and Shelf-life Characteristics". *J. Anim. Sci.*, 46 (1978), 658-665.
- [6] Huff, E.L. and Parrish, F.C.Jr. "Bovine Longissimus Muscle Tenderness as Affected by Postmortem Aging Time, Animal Age and Sex." *J. Food Sci.*, 58 (1993), 713-716.
- [7] Koochmaraie, M. "Biochemical Factors Regulating the Toughening and Tenderness Processes of Meat". *Meat Sci.*, 43 (1996), S193-S201.
- [8] Yousif, O.K. and Babiker, S.A. "The Desert Camel as a Meat Animal". *Meat Sci.*, 26 (1989), 245-254.
- [9] Shalash, M.R. *Provisional Report, International Foundation for Science*, 6, 1988.
- [10] Abouheif, M.A., Abdo, G.A., Basmaeil, S.M. and Al-Sobayel, A.A. "Identification of the Preference Patterns of Different Breeds of Sheep for Consumption in Saudi Arabia". *Asian-Australian J. Anim. Sci.*, 2 (1989), 129-132.
- [11] Al-Owaimer, N.A. "Changes in the Preferences of Saudi Citizens for Different Kinds of Meat and Sheep Breeds". *Alex.Sci.Exch.*, 20 (1999), 235-241.
- [12] El-Faer, M.Z., Rawdah, T.N., Attar, K.M. and Dawson, M.V. "Mineral and Proximate Composition of the Meat of the One-humped Camel (*Camelus Dromedarius*)". *Food Chem.*, 42 (1991), 139-143.
- [13] Elgasim, E.A. and Alkanhal, M.A. "Proximate Composition, Amino Acids and Inorganic Mineral Content of Arabian Camel Meat: Comparative Study". *Food Chem.*, 45 (1992), 1-4.
- [14] Dawood, A.A. "Physical and Sensory Characteristics of Najdi-Camel Meat". *Meat Sci.*, 39 (1995), 59-69.
- [15] Dawood, A.A. and Alkanhal, M.A. "Nutrient Composition of Najdi-Camel Meat". *Meat Sci.*, 39 (1995), 71-78.
- [16] Culler, R.D., Parrish, F.C. Jr., Smith, G.C. and Cross, H.R. "Relationship of Myofibril Fragmentation Index to Certain Chemical, Physical and Sensory Characteristics of Bovine Longissimus Muscles." *J. Food Sci.* 43 (1978), 1177-1180.
- [17] Cross, H.R., West, R.L. and Dutson, T.R. "Comparison of Methods for Measuring Sarcomere Length in Beef Semitendinosus Muscles." *Meat Sci.*, 5 (1981), 261-266.
- [18] SAS. *SAS User's Guide Statistics*. NC Cary: SAS Institute Inc., 1995.
- [19] Wheeler, T.L., Koochmaraie, M. and Crouse, J.D. "Effects of Calcium Chloride Injection and Hot-boning on the Tenderness of Round Muscles". *J. Anim. Sci.*, 69 (1991), 4871-4875.
- [20] Schmidt, G.R. and Keman, S. "Hot Boning and Vacuum Packaging of Eight Major Bovine Muscles". *J. Food Sci.*, 39 (1974), 140-142.
- [21] Koochmaraie, M., Whipple, G., Kretchmar, D.H., Crouse, J. D. and Mersmann, H.J. "Postmortem Proteolysis in Longissimus Muscle from Beef, Lamb, and Pork Carcasses". *J. Anim. Sci.*, 69 (1991), 617-624.
- [22] Morgan, J.B., Miller, K.R., Mendez, F.M., Hale, D.S. and Savell, J.W. "Using Calcium Chloride Injection to Improve Tenderness of Beef Mature Cows". *J. Anim. Sci.*, 69 (1991), 4469-4476.
- [23] Koochmaraie, M., Seideman, S.C., Schollmeyer, J.E., Dutson, T.R. and Crouse, J.D. "Effect of Post-mortem Storage on Ca Dependent Proteases, Their Inhibitor and Myofibril Fragmentation". *Meat Sci.*, 19 (1987), 187-196.
- [24] Taylor, R.G., Geesink, G.H., Thompson, V.F., Koochmaraie, M. and Goll D.G. "Is Z-disk Degradation Responsible for Postmortem Tenderization?" *J. Anim.Sci.*, 73 (1995), 1351-1367.
- [25] Moller, A.J. "Analysis of Warner-bratzler Shear Patterns with Regard to Myofibrillar and Connective Tissue Components of Tenderness". *Meat Sci.*, 5 (1981), 247-262.
- [26] Bailey, A.J. and Light, N.D. *Connective Tissue in Meat and Meat Products*. New York: Elsevier Applied Science, 1989.
- [27] Olson, D.G., Parrish, F.C. Jr. and Stromer, M.H. "Myofibril Fragmentation and Shear Resistance of three Bovine Muscle during Postmortem Storage". *J. Food Sci.*, 41 (1976), 1036-1041.
- [28] Wheeler, T.L., Savell, J.W., Cross, H.R., Hunt, D.K. and Smith, S.B. "Effect of Postmortem Treatments on the Tenderness of Meat from Hereford, Brahman and Brahman-cross Beef Cattle". *J. Anim. Sci.*, 68 (1990), 3677-3686.

- [29] Klotz, P.C., Baraco, V.M.H., Dijk, A. van, Smulders, F.J.M., Hoving-Bolink, A. H., Hulsegge, B. and Eikelenboom, G. "Effects of Rate of pH Fall, Time of Deboning, Aging Period, and Their Interaction on Veal Quality Characteristics". *J. Anim. Sci.*, 78 (2000), 1845-1851.
- [30] Herring, H.K., Cassens, R.C. and Briskey, E.J. "Further Studies on Quality Bovine Muscle Tenderness as Influenced by Dressing Position, Sarcomere Length and Fiber Diameter". *J. Food Sci.*, 30 (1965) 1049-1054.
- [31] Parrish, F.C. Jr., Young, R.B., Miner, B.E. and Anderson, L.D. "Effects of Postmortem Conditions on Calcium Chemical, Morphological and Organoleptic Properties of Bovine Muscle". *J. Food Sci.*, 38 (1973), 690-695.
- [32] Wheeler, T.L. and Koohmaraie, M. "Prerigor and Postrigor Changes in Tenderness of Ovine Longissimus Muscle". *J. Anim. Sci.*, 72 (1994), 1232-1238.

## تأثير فصل العضلة عن اللحم والتعتيق على فترات زمنية مختلفة على طراوة لحوم الإبل

عبد الله ناصر العويمر

قسم الإنتاج الحيواني، كلية الزراعة، جامعة الملك سعود،  
ص.ب. ٢٤٦٠ الرياض ١١٤٥١، المملكة العربية السعودية

( قدم للنشر في ١٣/١٠/١٤٢١ وقبل للنشر في ١٨/١٢/١٤٢١ هـ )

**ملخص البحث.** أجريت هذه التجربة لدراسة تأثير فصل العظم من اللحم بعد الذبح مباشرة والتعتيق على فترات زمنية مختلفة على طراوة لحوم الإبل. تم أخذ العضلة العينية ما بين الضلع ٧-١١ من كلا الجنسين من ٦ ذبائح من الإبل (متوسط وزن الذبيحة ١٣٣ كجم) ثم إزالة وتقطيع العضلة العينية إلى شرائح بسلك ٢،٥ سم بعد ساعة واحدة و٢٤ ساعة بعد الذبح، ومن ثم تغليفها تحت التفريغ الهوائي وتخزينها لمدة ١، ٣، ٧، ١٠، ١٤ يوم بعد الذبح على درجة حرارة ٢م. عند نهاية كل فترة تخزين، تم عمل القياسات التالية: قوة القطع للعضلة المطبوخة، درجة تكسير الليفة، طول الساركومير وفقد الرطوبة أثناء الطبخ والتخزين. أوضحت النتائج أن فصل العضلة بعد الذبح مباشرة زاد من قوة القطع معنوياً ( $P < 0.05$ ) في الأيام ٣ و٧ مقارنة بالعضلات المفصولة بعد ٢٤ ساعة من الذبح. وجد أن السبعة أيام الأولى من فترة التعتيق قللت بشكل واضح من قوة القطع أما في اليوم ١٠ و١٤ فكان التغيير قليل. كان معظم التغير في درجة تكسير الليفة خلال السبعة أيام الأولى بعد الذبح وأن فصل العضلة عن العظم بعد الذبح مباشرة قلل من درجة تكسير العضلة مقارنة بالفصل بعد ٢٤ ساعة. انخفض طول الساركومير معنوياً ( $P < 0.05$ ) عند فصل العضلة عن اللحم بعد ساعة واحدة من الذبح في يوم ٣ و٧ مقارنة بالفصل بعد ٢٤ ساعة. نستنتج من هذه الدراسة أن فصل اللحم عن العظم خلال ٢٤ ساعة الأولى من الذبح لا يؤثر على طراوة لحوم الإبل وأنه بالإمكان زيادة طراوة لحوم الإبل بالتعتيق لمدة ٧ أيام بعد الذبح.