

Tracing Monooxygenase Level by Synergistic Bioassays in Field and Sequential Generations of House Flies (Diptera: Muscidae)*

Deifalla H. Al-Rajhi

*Department of Plant Protection, College of Agriculture,
King Saud University, Riyadh, Saudi Arabia*

Abstract. Carbaryl, parathion, and diazinon bioassays were conducted in field through fourteen sequential generations of house fly, *Musca domestica* (L.) to trace monooxygenase level and role in the detoxification process. Parathion was more toxic to susceptible (S) strain (LC_{50} 0.013 μ g/vial) than diazinon (LC_{50} 0.33 μ g/vial). Carbaryl showed the lowest toxicity against the flies (LC_{50} 35 μ g/vial). The toxicities of carbaryl, parathion, and diazinon against S strains in relation to the field strains have been increased by 11, 45, and 346-folds respectively. Pretreatment of insects with piperonyl butoxide (Pb) inhibited the monooxygenase system, depriving the insects of a key defense mechanism against the three insecticides used in this study. Almost similar percent dependency values were obtained for the field strains (64, 62, and 61%) and S strains (71, 60, and 53%) on monooxygenases for detoxification of carbaryl, parathion, and diazinon respectively.

Introduction

Tolerance of insects to pesticides is a very known phenomenon. It can finally be due to any of several factors acting alone or in combination. In many cases, detoxification of insecticides by the metabolic action of enzymes is a major factor in insecticide tolerance. There is evidence, however, that monooxygenase detoxification mechanism is of far greater importance than others [1].

Biological assays are recommended for the detection and confirmation of insecticide tolerance or resistance. They can be conducted with insecticides and synergists to estimate relative detoxification role of the various enzyme systems. Brattsten and Metcalf [2,3,4], have suggested that the use of Pb which inhibits monooxygenases, could be used to alter the LD_{50} of carbaryl, which is metabolized by monooxygenases to indicate *in vivo* monooxygenase level in insects. Brindley [5] has indicated that synergist difference (SD) and percent dependency (%D) were more useful parameters in estimating monooxygenase or other enzyme system *in vivo* than synergist ratio.

Bioassays could effectively be used if the physiological and environmental factors are standardized [6]. Previous work, however, confirmed the effectiveness of the insecticide bioassay and revealed population differences in susceptibility and detoxification potential in field populations of black grass bugs [7] and pea aphids [8].

It is well established that monooxygenase activity shows considerable variation with respect to the physiological factors such as species, strain, sex, age, stage of development, and endogenous or exogenous factors and the environmental factors such as pesticide pressure, diet, and secondary plant substances [9,10,11].

Therefore, this study was organized to test the ability of the synergistic contact bioassay for tracing monooxygenase level in field through sequential generations of house fly and estimating its role in carbaryl, parathion, and diazinon detoxification. The data of the present study are interpreted by calculations of SD and %D values [5].

Materials and Methods

Insects

Laboratory strain of house flies was reared on natural diet of milk and sugar as described by Al-Rajhi *et al.* [12]. Pupae were collected from decayed matter and organic wastes located in the farm of Animal Production Department, Oleisha, Riyadh. Adults of the first generation were considered as a field strain, those of the fourteenth generation were considered susceptible strain.

Chemicals

In addition to the synergist, Pb, three insecticides were also used in this study: Carbaryl (1-naphthyl N-methylcarbamate (analytical grade, 99.4%). Parathion (0,0-diethyl -0-p-nitrophenyl phosphorothioate (analytical grade, 99.9%) and diazinon [0,0-diethyl-0- (2-isopropyl 4-methyl -6- pyrimidly) phosphorothiolate] (analytical grade, 99.9%). These chemicals were supplied by the United States Environmental Protection Agency.

Bioassay procedures

Chemicals and insects were introduced into the bioassay vials as follows:

Parathion, diazinon, carbaryl, and Pb were dissolved in acetone and diluted to make several concentrations of insecticides and 100 μg Pb per vial. A volume of 0.5 ml of each concentration was transferred to each vial. The solutions containing the

insecticides or the synergist were then allowed to evaporate by placing each vial side ways on a breadboard rolling the vials as evaporation occurred. This action permitted the solution inside the vials to leave a residue evenly distributed on the inner walls of the vials. Each bioassay included at least 5 different concentrations of insecticide and each concentration was replicated 4 times. The insects were pretreated with Pb by placing them in vials containing 100 μg residues of Pb. The insects were then transferred after 4h into vials containing several concentrations of insecticides. The vials were covered with cheesecloth. Five insects were used in each treated vial. The bioassay temperature was held at 25 ± 0.5 C. Several precautions were taken to insure that insects were alive and healthy during the 24h time interval of each experiment.

Results

Figures 1-3 show the susceptibility of house fly generations to carbaryl, parathion, and diazinon with and without Pb. The figures clearly indicate that the susceptibility of house fly generations to the three insecticides increased considerably as house flies advanced in generation. The toxicities of carbaryl, parathion, and diazinon have increased by 11, 45, and 346-fold respectively for the S strain compared to field strain (Fig. 4-6). Parathion was more toxic to the S strain (LC_{50} 0.013 μg per vial) than diazinon (LC_{50} 0.33 μg per vial). Carbaryl showed the lowest toxicity




Fig. 1. Susceptibility of house fly generations to carbaryl and carbaryl + piperonyl butoxide




Fig. 2. Susceptibility of house fly generations to parathion + and parathion + piperonyl butoxide




Fig. 3. Susceptibility of house fly generations to diazinon and diazinon + piperonyl butoxide




Fig. 4. Toxicity of carbaryl on susceptible strain and 3rd generation of house flies with and without piperonyl butoxide




Fig. 5. Toxicity of parathion on susceptible and field strains of house flies with and without piperonyl butoxide




Fig. 6. Toxicity of diazinon on susceptible and field strains of house flies with and without piperonyl butoxide

against the flies (LC_{50} 35 μg per vial). No mortality among field strain occurred in treatment with up to 2 mg carbaryl per vial, the highest carbaryl amount that could be dissolved in 0.5 ml acetone.

Table 1. Toxicity of synergized or unsynergized carbaryl, parathion, and diazinon to field and S strains of house fly.

Insecticide	Field strain			Susceptible strain		
	LC_{50} ($\mu\text{g/vial}$)	Synergized LC_{50}	%D on MFO	LC_{50} ($\mu\text{g/vial}$)	Synergized LC_{50}	%D on MFO
Carbaryl	400*	160	65	35	11	71
Parathion	4.5	1.7	63	0.015	0.005	63
Diazinon	15	6	62	0.44	0.23	47

*for the third generation

LC_{50} Values are calculated by liner regression

CSD is calculated from the equation: $\log \text{LC}_{50} = 1.014 \log (\text{CSD}) - 0.009$.

$$\%D = \frac{\text{Unsynergized } \text{LC}_{50} - \text{Synergized } \text{LC}_{50}}{\text{Calculated Synergist Difference (CSD)}}$$

Treatment with Pb enhanced the toxicity of carbaryl, parathion, and diazinon with similar degrees. The table shows the %D values of the flies on monoxygenase for detoxification of the three insecticides used in this study.

Discussion

The enhancement of carbaryl, parthion, and diazinon toxicities by Pb treatment revealed the relative importance of the flies monoxygenases in detoxification of the three insecticides. Metcalf and Fukuto [13] reported great enhancement of carbaryl toxicity against house flies when Pb was used.

Osman and Brindley [7] found that grass bug's monoxygenase detoxification was less important than that of the alfalfa leafcutting bee, where monoxygenases have an important role [14]. On the other hand, Al-Rajhi and Brindley [8] indicated that Pb did not enhance carbaryl toxicity against pea aphids. They concluded that aphids may be dependent on detoxification mechanism(s) other than monoxygenase.

The relative tolerance of the house fly field strains to carbaryl toxicity may be attributed to penetration or detoxification mechanism(s) other than monoxygenase.

The estimated %D values for monoxygenase detoxification of carbaryl, parathion, and diazinon were similar for the field and S strains indicating that environmental factors have little or no effect on flies monoxygenase level.

The data of this study confirmed the effectiveness of the synergistic bioassays in revealing monoxygenase level and role in house flies and obviate the possible alternative detoxification mechanism (s). It also indicated that environmental factors have little or no effect on monoxygenase level in house flies, under such limited and homogeneous population of insect inhabiting the location.

Acknowledgement. This work was supported by the Agricultural Research Center, College of Agriculture, King Saud University.

References

- [1] Georghiou, G.P. "The Evolution of Resistance to Pesticides." *Ann. Rev. Ecol. yst.* 3(1972), 133-168.
- [2] Brattsten, L.B. and Metcalf, R.I.L. "The Synergistic Ratio of Carbaryl and Piperonyl Butoxide as an Indicator of the Distribution of Multi-function Oxidases in Insecta." *J. Econ. Entomol.*, 63 (1970), 101-104.
- [3] Brattsten, L.B. and Metcalf, R.I.L. "Age-dependent Variations in the Response of Several Species of Diptera to Insecticidal Chemicals." *Pestic. Biochem. Physiol.*, 3(1973a), 189-198.

- [4] Brattsten, L.B. and Metcalf, R.L. "Synergism of Carbaryl Toxicity in Natural Insect Populations." *J. Econ. Entomol.*, 68 (1973b), 227-230.
- [5] Brindley, W.A. "Synergist Differences as an Alternate Interpretation of Carbaryl Piperonyl Butoxide Toxicity Data." *Environ. Entomol.*, 6(1977), 885-888.
- [6] Brindley, W.A. "Insecticidal Bioassays with Field Populations of Alfalfa Weevils, a Simplified Approach." *J. Econ. Entomol.*, 68(1975), 227-230.
- [7] Osman, D.H. and Brindley, W.A. "Estimating Monooxygenase Detoxification in Field Populations: Toxicity and Distribution of Carbaryl in Three Species of Labops Grassbugs." *Environ. Entomol.*, 10 (1981), 676-680.
- [8] Al-Rajhi, D.H. and Brindley, W.A. "Relationships between Synergistic and Biochemical Analysis of Paraoxon Tolerance in Pea Aphids." *Alex. Sci. Exch.*, 6(1985), 55-72.
- [9] Wilkinson, C.F. *Insecticide Biochemistry and Physiology*. New York: Plenum Press, 1976.
- [10] Brattsten, L.B., C.F., Wilkinson, and Eisner, T. "Herbivore Plant Interactions: Mixed Function Oxidases and Secondary Plant Substances." *Science*, 196 (1977), 1349-1381.
- [11] Walker, C.A. "Species Differences in MFOs Activity and Their Relationship to Biological Half-Lives." *Drug Metab. Rev.*, 7 (1978), 295-323.
- [12] Al-Rajhi, D.H., Moussa, M.E., and Morshedy, M. "Insecticidal Susceptibility on Two Strains of House Fly, *Musca domestica* L. in Saudi Arabia." *J. Coll. Agric. King Saud Univ.*, 7 (1985), 467-474.
- [13] Metcalf, R.L. and Fukuto, T.R. "Effects of Chemical Structure on Intoxication and Detoxication of Phenyl N-methyl Carbamates in Insects." *J. Agric. Food. Chem.*, (1965), 220.
- [14] Guirguis, G.N. and Brindley, W.A. "Carbaryl Penetration into and Metabolism by Alfalfa Leafcutting Bees, *Megachile pacifica*." *J. Agric. Food Chem.*, 23 (1975), 274-279.

الكشف عن مستوى النشاط الأنزيمي للمونوأكسجينيز بواسطة اختبارات التقييم الحيوي المنشط على سلالات حقلية وحساسة من الذباب المنزلي

ضيف الله هادي الراجحي

قسم وقاية النبات، كلية الزراعة، جامعة الملك سعود

الرياض، المملكة العربية السعودية

ملخص البحث. أجريت اختبارات التقييم الحيوي للمبيدات: كرابريل - باراثيون، ديازينون على سلالة حقلية وعلى مدى أربعة عشر جيلاً متتابعاً من سلالة معملية من الذباب المنزلي *Musca domestica* (L)، لقياس مستوى النشاط الأنزيمي للمونوأكسجينيز، وكذلك لبيان دوره في عمليات تقليل سمية المبيدات المختبرية. وقد أظهرت النتائج أن الباراثيون كان أكثر سمية للسلالة الحساسة من الذباب المختبر حيث كان التركيز القاتل النصفى (ت. ق. ٥٠) من الباراثيون ٠,١٣، ٠ ميكروجراماً/أنبوبة بينما كانت قيمة (ت. ق. ٥٠) للديازينون ٠,٣٣، ٠ ميكروجراماً/أنبوبة أما الكرابريل فقد كان أضعف المبيدات الثلاثة في هذا المجال حيث كانت قيمة ت. ق. ٥٠ له ٠,٣٥، ٠ ميكروجراماً/أنبوبة. كما أظهرت النتائج أيضاً أن سمية كل من كرابريل وباراثيون وديازينون للسلالة الحساسة (المعملية) قد تضاءلت عن سميتها بمقدار ١١، ٤٥، ٣٤٦ ضعفاً على التوالي.

معاملة الحشرات بالمثبط بيرونيل بيوتوكسايد أدت إلى تثبيط إنزيمات الـ مونوأكسجينيز حيث وجد أن نسب اعتماد السلالة الحقلية والحساسة على هذه الإنزيمات في تقليل سمية المبيدات الثلاثة تكاد تكون متقاربة حيث كانت ٦٤٪، ٦٢٪، ٦١٪ للسلالة الحقلية و ٧١٪، ٦٠٪، ٥٣٪ للسلالة الحساسة لكل من الكرابريل والباراثيون والديازينون على التوالي.

